



THE HUMAN FAMILY  
RESEARCH CENTER

*Every Human Being  
Leaves a Fingerprint on the Universe*

# ***FACTS IN FOCUS***

## **Toward an Understanding of Why Induced Abortion Can Be More Traumatizing than Spontaneous Abortion**

---

Unlike the pain of spontaneous abortion (miscarriage) which tends to lessen over time, the pain of abortion may worsen in the months and years following the procedure [1-2].

There are a number of reasons why induced abortion can lead to more severe and insidious psychological distress than miscarriage [3].

1. Miscarriage is an involuntary loss. Even so, some women blame themselves for their loss insisting they could have done something differently to prevent it from happening. Abortion however is voluntary. Thus, many women experience guilt, self-criticism, or even self-loathing [4,5,6, 14].
2. Sadness and grief are culturally recognized with miscarriage; whereas grief after induced abortion is not. As a result, women may suppress abortion-related emotions, as Kluger-Bell, a psychotherapist stated: *“When other people are reluctant to listen to us, when there are no ceremonies to publicly acknowledge the impact of our experiences, we receive the covert message that others would rather not hear what we have to say, and this makes it difficult to even identify our reactions to our losses.”* [7]
3. The grief process after abortion is likely to be more complicated than with miscarriage because grief resolution requires acknowledgment of loss, in this case a death. Rather than acknowledging the existence of their fetal child and/or their role in his/her death,, women who have aborted may find themselves either in denial and avoidance or having unwanted reexperiencing and intrusive feelings, neither of which they can successfully control. The traumagenic nature of abortion and how this complicates grief resolution has been previously reported. [15] Unhealthy coping mechanisms can include escalating dependencies on alcohol, drugs, food, work and sex which can in turn increase symptoms of anxiety and depression.
4. The available peer reviewed scientific research indicates that adverse mental health effects of induced abortion persist longer than the negative effects of miscarriage [9- 13].  
Repression of grief, lack of support, self-recrimination, exacerbation or development of anxiety or depression, and/or abuse of substances are all factors that may render voluntary perinatal loss potentially more difficult to move beyond when compared to involuntary perinatal loss.

- 
1. Bradley CF: Abortion and subsequent pregnancy, *Canadian Journal of Psychiatry* 1984; 29, p 494.
  2. Devore NE. The relationship between previous elective abortions and postpartum depressive reactions. *Journal of Obstetric Gynecologic and Neonatal Nursing* 1979; July/August: 237-240.
  3. Coleman PK. (in press). The Psychological Pain of Perinatal Loss and Subsequent Parenting Risks: Could Induced Abortion Be More Problematic than Other Forms of Loss? *Current Women’s Health Reviews*.
  4. Kero A, Hoegberg U, Jacobsson L, Lalos A. Legal abortion: A painful necessity. *Social Science & Medicine* 2001; 53: 1481-1490.
  5. Miller WB, Pasta DJ, Dean CL. Testing a model of the psychological consequences of abortion. In Beckman LJ and Harvey SM, Eds. *The new civil war: The psychology, culture, and politics of abortion*. Washington, DC: American Psychological Association 1998: 235-267.

6. Skelton G. Many in survey who had abortions cite guilt feelings. Los Angeles Times. (March 19, 1989).
7. Kluger-Bell K. *Unspeakable losses: Healing from spontaneous abortion, abortion, and other pregnancy losses* 2000; New York, HarperCollins Publishers.
8. Speckhard AC, Rue VM. Complicated mourning: Dynamics of impacted post-abortion grief. *Pre-and Perinatal Psychology Journal* 1993; 8: 5-32.
9. Peppers LG, Knapp RJ. Maternal reactions to involuntary fetal/infant death. *Psychiatry* 1980; 43: 155-159.
10. Broen AN, Moum T, Bodtke AS, Ekeberg O. Psychological impact on women of spontaneous abortion versus induced abortion: A 2-year follow-up study. *Psychosomatic Medicine* 2004; 66, 265-271.
11. Broen AN, Moum T, Bødtker AS, Ekeberg O. The course of mental health after miscarriage and induced abortion: a longitudinal, five-year follow-up study. *BMC Medicine* 2005; 12, 18.
12. Broen AN, Moum T, Bødtker AS, Ekeberg O. Predictors of anxiety and depression following pregnancy termination: a longitudinal five-year follow-up study. *Acta Obstet Gynecol Scand.* 2006; 85:317-23.
13. Coleman PK, Reardon DC, Cogle J. Substance use among pregnant women in the context of previous reproductive loss and desire for current pregnancy. *British Journal of Health Psychology* 2005; 10: 255-268.
14. Rue VM, Coleman PK, Rue JJ, Reardon, DC. Induced abortion and traumatic stress: A preliminary comparison of American and Russian women. *Medical Science Monitor* 2004; 10, SR5-16.